



2016 Annual and Perennial Ryegrass and Festulolium Report

G.L. Olson, S.R. Smith, and T.D. Phillips, Plant and Soil Sciences

Introduction

Annual ryegrass (*Lolium multiflorum*) and perennial ryegrass (*Lolium perenne*) are high-quality, productive, cool-season grasses used in Kentucky. Both have exceptionally high seedling vigor and are highly palatable to livestock. In Kentucky, winter survival can be an issue for many annual ryegrass varieties, so before planting, review winter survival results in this publication. The severe winter of 2013-14 showed those varieties that are not adapted to Kentucky.

Annual ryegrasses are increasing in use across Kentucky as more winter-hardy varieties are released and promoted. Annual ryegrass is productive for three to four months and is used primarily for late fall and early to late spring pasture. Winter growth occurs only during mild winters. This crop has garnered increased interest for high-quality baleage. Two main types of annual ryegrasses are used. The most commonly used type in Kentucky is Italian ryegrass. The other is sometimes referred to as Westerwolds ryegrass. The Westerwolds type is a true annual, in that stands seeded in the spring produce seedheads that summer, and little regrowth occurs after seedheads are produced. Westerwolds ryegrass varieties are commonly used in the lower South (Florida to Texas) because they can be seeded in the fall and will survive the winter. Many varieties also survive Kentucky

winters. Italian ryegrass is native to Southern Europe and is not a true annual. Italian ryegrasses provide high yields of quality forage and show quick regrowth. If planted in the spring, no or few seedheads will grow that summer (vernalization is required). Spring planting of Italian ryegrass is common in northern states (e.g., Wisconsin, Minnesota, etc.) for summer grazing, but most current varieties do not dependably survive Kentucky summers. Italian ryegrasses are almost always planted late summer to early fall in Kentucky and typically provide forage production into early summer, often one to two months later than Westerwolds types.

Perennial ryegrass can be used as a short-lived hay or pasture plant and has growth characteristics similar to tall fescue. It is more persistent than Italian ryegrass but less persistent than other cool-season grass species. It tillers more profusely but is lower growing than Italian ryegrass and will not form a seedhead in the seeding year. Both diploid (two sets of chromosomes) and tetraploid (four sets of chromosomes) varieties of perennial ryegrass exist. Tetraploids have larger tillers and seedheads and wider leaves. Tetraploid types tend to be

taller and less dense than diploid types even in early stages of regrowth. Diploid types produce more tillers, have better stand persistence, and are more tolerant to heavy grazing.

Intermediate or hybrid ryegrass (*Lolium hybridum*, Hausska) is the result of a cross between Italian ryegrass and perennial ryegrass. It is not as winter hardy as perennial ryegrass, but it is higher yielding. It is also more persistent and winter hardy than Italian ryegrass. Its uses are similar to those of perennial ryegrass but typically only survive two years or less in Kentucky.

Both forage and turf types of annual and perennial ryegrasses are available. Turf types are low growing and have poor yield. Turf types are also infected with a fungal endophyte that lives inside the plant, protecting it from insect attack but producing a toxin that reduces performance of grazing animals. All turf types are infected. Plant only forage-type varieties for grazing, hay, or silage.

Festuloliums are hybrids between various fescues and ryegrasses with higher quality than tall fescue and improved stand survival over perennial ryegrass. Their use in Kentucky is still limited since they do not survive as long as tall fescue

Table 1. Temperature and rainfall at Lexington, Kentucky, in 2013, 2014, 2015, and 2016.

	2013				2014				2015				2016			
	Temp		Rainfall		Temp		Rainfall		Temp		Rainfall		Temp		Rainfall	
	°F	DEP ¹	IN	DEP	°F	DEP	IN	DEP	°F	DEP	IN	DEP	°F	DEP	IN	DEP
JAN	38	+7	4.50	+1.64	25	-6	2.28	-.58	32	+1	2.17	-0.69	32	+1	0.80	-2.06
FEB	36	+1	1.78	-1.43	30	-5	5.47	+2.26	26	14	3.08	-0.13	38	+3	6.09	+2.88
MAR	39	-5	5.47	+1.07	39	-5	3.08	-1.32	45	+1	7.34	+2.94	52	+8	4.07	-0.33
APR	55	0	4.46	+0.58	58	+3	5.27	-1.89	57	+2	13.19	+9.31	57	+2	3.97	+0.09
MAY	65	+1	5.23	+0.76	66	+2	5.72	+1.25	69	+5	3.02	-1.45	64	0	9.17	+4.70
JUN	72	0	7.32	+3.66	75	+3	2.93	-0.73	75	+3	8.20	+4.54	76	+4	5.09	+1.43
JUL	72	-4	9.33	+4.33	74	-2	3.18	-1.82	77	+1	10.22	+5.22	79	+3	7.43	+2.43
AUG	72	-3	3.68	-0.25	76	+1	6.53	+2.60	74	-1	3.49	-0.44	79	+4	4.37	+0.44
SEP	67	-1	2.21	-0.99	69	+1	3.63	+4.3	72	+4	3.49	+0.29	74	+6	2.18	-1.02
OCT	55	-2	7.02	+4.45	57	0	5.55	+2.98	59	+2	2.78	+0.21	64	+7	0.37	-2.20
NOV	41	-4	3.06	-0.33	41	-4	2.79	-0.60	51	+6	3.72	+0.33				
DEC	36	0	4.19	+0.21	40	+4	2.47	-1.51	49	+13	8.42	+4.44				
Total			58.25	+13.70			49.4	+4.85			69.12	+24.57			43.54	+6.36

¹ DEP is departure from the long-term average.

² 2016 data is for the ten months through October.

but some of the newer varieties are more adapted to Kentucky environmental conditions.

This report provides current yield data on annual and perennial ryegrass varieties in trials in Kentucky as well as guidelines for selecting varieties. Tables 16, 17, and 18 show summaries of all annual and perennial ryegrass and festulolium varieties tested in Kentucky for the last 15 years. The UK Forage Extension website at www.uky.edu/Ag/Forage contains electronic versions of all forage variety testing reports from Kentucky and surrounding states and a large number of other forage publications.

Important Selection Considerations

Local adaptation and seasonal yield. The variety should be adapted to Kentucky as indicated by good winter survival and good performance across years and locations in replicated yield trials, such as those presented in this publication. Choose high-yielding varieties, but choose varieties that are productive during the desired season of use.

Seed quality. Buy premium-quality seed that is high in germination, high in purity, and free from weed seed. Buy certified seed or proprietary seed of an improved variety. An improved variety is one that has performed well in independent trials. Other information on the label will include the test date (which must be within the previous nine months), the level of germination, and percentage of other crop and weed seed. Order seed well in advance of planting time to assure that it will be available when needed.

Important: When seeding perennial ryegrasses for horse or cattle pastures (of any kind), insist on an endophyte-free variety. The endophyte level will be stated on a green tag on every bag of seed. Most forage types of perennial ryegrass are endophyte free, and most new turf types are infected. This endophyte is similar to the endophyte of tall fescue and produces alkaloids that are toxic to cattle and horses.

Description of the Tests

Data from eleven studies are reported. Annual ryegrass tests were established in the fall of 2012, 2013, 2014, and 2015

Table 2. Descriptive scheme for the stages of development in perennial forage grasses.

Code	Description	Remarks
Leaf development		
11	First leaf unfolded	Applicable to regrowth of established (plants) and to primary growth of seedlings.
12	2 leaves unfolded	Further subdivision by means of leaf development index (see text).
13	3 leaves unfolded	
•	•••••	
19	9 or more leaves unfolded	
Sheath elongation		
20	No elongated sheath	Denotes first phase of new spring growth after overwintering. This character is used instead of tillering which is difficult to record in established stands.
21	1 elongated sheath	
22	2 elongated sheaths	
23	3 elongated sheaths	
•	•••••	
29	9 or more elongated sheaths	
Tillering (alternative to sheath elongation)		
21	Main shoot only	Applicable to primary growth of seedlings or to single tiller transplants.
22	Main shoot and 1 tiller	
23	Main shoot and 2 tillers	
24	Main shoot and 3 tillers	
•	•••••	
29	Main shoot and 9 or more tillers	
Stem elongation		
31	First node palpable	More precisely an accumulation of nodes. Fertile and sterile tillers distinguishable.
32	Second node palpable	
33	Third node palpable	
34	Fourth node palpable	
35	Fifth node palpable	
37	Flag leaf just visible	
39	Flag leaf ligule/collar just visible	
Booting		
45	Boot swollen	
Inflorescence emergence		
50	Upper 1 to 2 cm of inflorescence visible	
52	¼ of inflorescence emerged	
54	½ of inflorescence emerged	
56	¾ of inflorescence emerged	
58	Base of inflorescence just visible	
Anthesis		
60	Preanthesis	Inflorescence-bearing internode is visible. No anthers are visible.
62	Beginning of anthesis	First anthers appear.
64	Maximum anthesis	Maximum pollen shedding.
66	End of anthesis	No more pollen shedding.
Seed ripening		
75	Endosperm milky	Inflorescence green
85	Endosperm soft doughy	No seeds loosening when inflorescence is hit on palm.
87	Endosperm hard doughy	Inflorescence losing chlorophyll; a few seeds loosening when inflorescence hit on palm
91	Endosperm hard	Inflorescence-bearing internode losing chlorophyll; seeds loosening in quantity when inflorescence hit on palm.
93	Endosperm hard and dry	Final stage of seed development; most seeds shed.

Source: J. Allan Smith and Virgil W. Hayes. 14th International Grasslands Conference Proc. p. 416-418. June 14-24, 1981, Lexington, Kentucky.

at Lexington. Perennial ryegrass tests and festulolium tests were established in 2013, 2014, and 2015 at Lexington. The soil at Lexington is a well-drained silt loam (Maury) and is well suited for ryegrass production.

Seedlings were made at the rate of 25 pounds per acre into a prepared seedbed with a disk drill. Plots were 5 feet by 20

feet in a randomized complete block design with four replications with a harvested plot area of 5 feet by 15 feet. For the perennial tests nitrogen was top-dressed at 60 pounds per acre of actual nitrogen in March, May, and August. For the annual tests nitrogen was top-dressed at 60 pounds per acre in March and 60 pounds after the first spring harvest. The tests

Table 3. Dry matter yields, seedling vigor, plant height, maturity, and stand persistence of annual ryegrass varieties sown August 31, 2012, at Lexington, Kentucky (see Table 16 for designation of Italian or Westerwolds type commercial varieties).

Variety	Seedling Vigor ¹ Oct 11, 2012	Maturity ²					Percent Stand					Height (inches)					Yield (tons/acre)				
		2013					2012					2013					2012				
		Apr 22	May 21	Jun 11	Jun 25	Oct 11	Mar 20	Jul 22	Aug 21	Dec 14	Apr 22	May 21	Jun 11	Jun 25	Dec 14	Apr 23	May 21	Jun 11	Jun 25	Jul 23	Total
Commercial Varieties-Available for Farm Use																					
MX108(Max)	4.6	32.0	47.3	56.0	56.5	100	100	100	100	8	14	19	18	13	0.95	1.82	1.25	0.60	0.22	0.45	5.30*
TetraPrime	3.5	31.8	46.8	57.0	57.0	100	100	100	100	6	13	18	15	13	0.51	1.78	1.34	0.49	0.23	0.37	4.72*
LHT-102	4.0	31.5	49.3	53.0	56.0	100	100	100	100	6	12	19	10	11	0.44	1.77	1.55	0.39	0.22	0.31	4.67
Marshall	4.6	32.8	50.8	59.5	62.0	100	100	20	4	9	16	21	14	7	1.28	1.90	0.99	0.44	0.05	0.00	4.66
AE110	3.4	32.5	50.8	56.5	61.0	100	94	89	56	8	15	21	16	12	0.85	1.59	1.28	0.51	0.11	0.31	4.64
Centurion	4.0	33.3	46.3	60.0	62.0	100	100	30	8	9	17	17	9	9	0.99	2.12	0.86	0.46	0.09	0.00	4.53
TAMTBO	4.1	32.8	55.5	57.5	62.0	100	93	25	7	7	14	22	13	7	0.97	1.69	1.36	0.38	0.05	0.00	4.44
Feast II	4.9	31.3	53.5	55.0	56.0	100	70	93	92	8	9	24	13	13	0.96	0.98	1.41	0.49	0.20	0.31	4.35
Winterhawk	4.8	32.5	49.3	62.0	62.0	100	100	24	10	9	17	16	16	7	1.09	1.82	0.94	0.37	0.06	0.00	4.29
Jackson	4.1	32.8	52.0	62.0	62.0	100	100	13	4	10	16	20	14	8	1.20	1.69	0.98	0.36	0.02	0.00	4.24
TillageMaxBristol ³	2.3	32.3	54.0	57.5	62.0	89	87	18	4	7	14	21	12	9	0.98	1.60	1.20	0.38	0.08	0.00	4.24
TillageMaxINDY ³	2.1	32.3	51.3	59.5	62.0	91	90	14	3	6	15	21	14	10	0.76	1.74	1.20	0.41	0.07	0.00	4.19
TillageRootMax	2.3	33.0	48.8	60.0	62.0	98	100	13	1	7	17	19	15	9	0.65	1.97	1.07	0.42	0.07	0.00	4.18
Fria	4.9	33.3	50.8	62.0	62.0	100	100	20	1	9	16	17	15	8	1.09	1.66	0.92	0.41	0.08	0.00	4.16
Bruiser	5.0	32.5	51.3	61.0	62.0	100	100	23	5	9	15	17	14	8	1.24	1.51	0.93	0.29	0.05	0.00	4.02
Gulf	4.8	31.3	62.0	59.0	62.0	100	63	6	1	8	9	28	9	7	1.06	0.76	1.26	0.19	0.07	0.00	3.34
Experimental Varieties																					
PS-Lm-09-2	4.6	33.3	49.3	57.0	56.5	100	100	100	97	8	17	18	18	11	0.82	2.00	1.12	0.64	0.15	0.52	5.25*
Lh 4x-1PS	3.8	32.3	53.5	57.0	59.0	100	100	94	91	8	14	20	17	13	0.74	1.83	1.34	0.53	0.18	0.37	4.98*
Amp	3.1	33.0	56.0	60.0	62.0	99	98	30	11	7	17	21	15	8	0.79	1.80	1.29	0.47	0.08	0.00	4.43
IS-LWT 12	4.0	32.5	57.0	59.5	62.0	100	85	31	18	8	14	24	14	8	1.05	1.44	1.22	0.41	0.05	0.00	4.17
IS-LWT 14	3.6	32.0	55.5	58.5	62.0	100	99	36	16	8	14	21	15	8	0.86	1.54	1.24	0.48	0.06	0.00	4.17
IS-LWT 13	3.8	32.8	56.0	57.0	62.0	93	84	35	14	7	13	23	15	8	0.78	1.50	1.30	0.47	0.07	0.00	4.13
Mean	3.9	32.4	52.1	58.5	60.5	99	94	46	34	8	14	20	14	9	0.91	1.66	1.18	0.44	0.10	0.12	4.41
CV,%	15.2	2.1	5.6	3.5	1.4	4	12	20	23	13	10	10	13	13	22.83	12.34	12.54	22.47	28.01	44.11	9.99
LSD,0.05	0.9	0.9	4.1	2.9	1.2	5	16	13	11	1	2	3	3	2	0.29	0.29	0.21	0.14	0.04	0.07	0.62

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

³ These are mixtures that included crimson clover and/or tillage radish.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 4. Dry matter yields, seedling vigor, winter injury, plant height, maturity, and plant persistence of annual ryegrass varieties sown September 5, 2013, at Lexington, Kentucky (see Table 16 for designation of Italian or Westerwolds type commercial varieties).

Variety	Seedling Vigor ¹ Oct 14, 2013	Percent Stand		Winter Injury ² Jan 27	Plant Height(in) May 1	Maturity ³			Yield (tons/acre)				
		2013	2014			2014			2014				
		Oct 14	Apr 2			May 1	May 22	Jun10	May 2	May 23	Jun 10	Jun 27	Total
Commercial Varieties-Available for Farm Use													
Fria	4.3	96	97	1.5	19	37	50	60	2.01	0.89	0.49	0.09	3.47*
Ed	4.1	97	96	2.8	19	41	51	59	1.77	0.93	0.55	0.11	3.35*
Bruiser	4.8	97	96	2.5	19	42	53	58	1.80	0.88	0.55	0.12	3.34*
Marshall	3.8	93	95	1.3	22	36	48	59	1.94	0.82	0.47	0.10	3.33*
Barmultra II	3.4	89	70	3.5	15	33	52	56	1.20	1.05	0.72	0.18	3.15*
Assist	3.5	93	93	3.0	18	39	49	60	1.57	0.79	0.50	0.07	2.92*
Amp	3.4	89	46	3.5	17	43	54	59	1.09	0.75	0.54	0.11	2.49
Dyna-Gain	3.8	93	83	3.5	19	39	51	59	1.13	0.66	0.46	0.10	2.35
Jackson	4.0	95	78	5.3	16	49	54	61	1.10	0.68	0.43	0.10	2.31
Hercules	4.3	92	44	5.8	14	34	52	59	0.88	0.71	0.52	0.17	2.28
Nelson	3.3	90	51	4.8	15	34	53	60	0.91	0.64	0.51	0.10	2.16
Attain	3.6	92	31	4.5	14	33	53	61	0.71	0.58	0.36	0.09	1.74
Feast II	3.3	88	12	8.3	10	32	54	59	0.40	0.42	0.53	0.23	1.58
Verdure	4.0	92	23	7.5	12	32	53	60	0.51	0.46	0.38	0.06	1.41
Big Boss	3.3	90	13	7.3	12	34	56	60	0.43	0.39	0.38	0.07	1.27
Gulf	4.0	93	14	7.3	11	38	56	60	0.33	0.28	0.25	0.04	0.90
Dixie Gold	2.8	80	2	8.0	11	33	55	60	0.18	0.17	0.23	0.05	0.64
Experimental Varieties													
M2CVS	3.4	93	97	1.5	20	39	50	59	1.99	0.81	0.46	0.14	3.40*
ME4	3.3	88	89	1.3	23	37	48	58	1.98	0.83	0.52	0.04	3.37*
ME-94	3.9	92	95	2.0	20	42	50	58	1.75	0.86	0.49	0.10	3.19*
LMT-15M3	3.4	80	75	3.3	16	34	51	57	1.30	0.92	0.65	0.14	3.01*
B-13.0171	2.8	84	21	5.0	15	34	50	61	0.78	0.54	0.60	0.12	2.03
FL4XMep	2.8	80	20	4.8	14	46	55	62	0.64	0.55	0.38	0.10	1.67
FL4XMarmi	2.6	70	27	5.3	15	44	56	61	0.50	0.34	0.32	0.11	1.26
FL4XMaron	2.8	84	9	4.8	12	47	56	62	0.38	0.23	0.31	0.07	0.99
Mean	3.5	89	55	4.3	16	38	52	59	1.09	0.65	0.46	0.10	2.30
CV,%	15.9	11	26	25.8	10	13	5	3	23.00	24.17	24.22	49.33	20.15
LSD,0.05	0.8	14	21	1.6	2	7	4	3	0.35	0.24	0.16	0.07	0.65

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Winter injury score based on a scale of 1 to 9 with 9 being the greatest amount of injury.

³ Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

were harvested using a sickle-type forage plot harvester. The first cutting was harvested at each location when all ryegrass varieties had reached at least the boot stage. Fresh weight samples were taken at each harvest to calculate dry matter production. Management practices for these tests regarding establishment, fertility (P, K, and lime are based on regular soil tests), weed control, and harvest timing were in accordance with University of Kentucky recommendations.

Results and Discussion

Weather data for Lexington are presented in Table 1.

Ratings for maturity (see Table 2 for maturity scale) and dry matter yields (tons/A) are reported in tables 3 through 12. Yields are given by cutting date for 2016 and as total annual production. Stated yields are adjusted for percent

weeds; therefore, the tonnage given is for crop only. Varieties are listed by total yield in descending order. Experimental varieties, listed separately at the bottom of the tables, are not available commercially.

In most years, annual ryegrasses can be expected to die or become unproductive after mid-June in their first summer. Unlike annual ryegrasses, perennials should be productive under Kentucky conditions for an average of two to three growing seasons.

Statistical analyses were performed on all data (including experimentals) to determine if the apparent differences are truly due to varietal differences or just due to chance. Varieties not significantly different from the top variety in the column are marked with one asterisk (*). To determine if two varieties are truly different, compare the difference between

them to the Least Significant Difference (LSD) at the bottom of the column. If the difference is equal to or greater than the LSD, the varieties are truly different when grown under the conditions at the given locations. The Coefficient of Variation (CV) is a measure of the variability of the data and is included for each column of means. Low variability is desirable; increased variability within a study results in higher CVs and larger LSDs.

Tables 13, 14, and 15 summarize information about distributors and yield performance for all annual and perennial ryegrass and festulolium varieties currently included in tests discussed in this report. Varieties are listed in alphabetical order by species, with the experimental varieties at the bottom. Remember that experimental varieties are not available for farm use; commercial varieties can be purchased from agricultural distributors.

Table 5. Dry matter yields, seedling vigor, and stand persistence of annual ryegrass varieties sown September 5, 2014, at Lexington, Kentucky (see Table 16 for designation of Italian or Westerwolds type commercial varieties).

Variety	Seedling Vigor ¹ Oct 9, 2014	Percent Stand		Winter Injury ² Jan 19,2015	Maturity ³		Plant Height(in) Apr 23	Yield (tons/acre)				
		2014	2015		2015			2014	2015			
		Oct 9	Apr 3		Apr 23	May 19		Dec 15	Apr 24	May 19	Jun 15	Total
Commercial Varieties- Available for Farm Use												
Centurion	4.1	99	95	0.5	32.3	50.5	18	0.53	1.51	1.22	0.48	3.74*
Winterhawk	4.1	99	92	1.5	35.8	54.5	18	0.74	1.29	1.05	0.31	3.39*
Bruiser	4.4	99	80	2.5	32.5	55.5	17	0.74	0.98	0.90	0.36	2.98
Ed	3.3	94	70	2.0	31.5	55.5	13	0.45	0.88	1.13	0.39	2.85
Marshall	4.0	98	81	0.5	32.5	54.0	17	0.51	1.14	0.91	0.28	2.84
Jackson	3.8	98	88	1.0	32.5	55.5	16	0.57	1.18	0.81	0.29	2.84
TetraPrime	3.0	98	98	0.5	31.0	46.3	12	0.34	1.17	0.99	0.22	2.72
Kowinearly	3.1	97	91	1.8	41.3	59.0	17	0.30	1.14	0.95	0.31	2.70
Meroa	3.0	78	64	2.8	36.3	55.0	13	0.28	0.96	0.93	0.48	2.65
Green Farm	4.0	100	66	2.8	46.0	59.5	21	0.52	0.87	0.71	0.29	2.40
Fria	4.0	99	58	3.3	32.0	57.0	13	0.44	0.58	0.96	0.32	2.30
Kospeed	4.4	99	63	1.8	41.0	59.0	16	0.58	0.72	0.78	0.19	2.26
TAMTBO	2.9	78	43	1.8	31.5	57.5	13	0.30	0.63	1.04	0.26	2.23
Tam 90	3.8	100	26	5.0	30.0	58.0	10	0.64	0.34	0.92	0.32	2.22
Nelson	2.1	65	48	1.8	31.0	56.5	12	0.34	0.70	0.90	0.26	2.20
Big Boss	2.5	68	6	5.8	30.3	58.0	11	0.48	0.21	0.86	0.52	2.07
Attain	2.1	63	16	2.3	30.5	58.0	11	0.38	0.32	0.92	0.32	1.95
Gulf	4.0	100	26	6.3	30.8	59.0	10	0.53	0.31	0.76	0.34	1.95
Big Bang	3.4	86	25	4.3	30.0	55.5	9	0.28	0.30	0.92	0.41	1.91
Bill	3.6	88	11	5.0	30.3	59.0	9	0.43	0.26	0.76	0.29	1.75
Verdure	3.4	87	4	6.0	29.5	59.5	9	0.51	0.18	0.72	0.26	1.66
Feast II	3.0	88	13	7.3	35.0	54.0	9	0.35	0.24	0.73	0.26	1.58
Experimental Varieties												
ME94	3.9	92	88	0.5	36.8	54.0	18	0.60	1.33	1.09	0.54	3.57*
ME4	2.9	96	94	0.0	32.3	53.0	19	0.43	1.40	0.91	0.41	3.14*
M2CVS	3.9	99	92	0.5	31.5	54.0	14	0.50	1.31	0.95	0.35	3.12*
GO-ITT12	3.0	94	89	1.8	32.5	55.0	15	0.28	0.93	1.28	0.29	2.79
GO-FLN2	3.3	97	49	3.3	30.5	57.0	10	0.50	0.49	1.08	0.32	2.40
GO-IT213	2.3	78	11	5.3	30.8	59.5	13	0.27	0.23	0.74	0.36	1.60
Mean	3.4	90	57	2.8	33.0	56.0	14	0.46	0.77	0.93	0.34	2.49
CV,%	20.9	9	36	33.2	14.0	3.0	18	30.34	32.57	24.47	47.53	17.71
LSD,0.05	1.0	12	21	1.3	7.0	2.0	4	0.20	0.35	0.32	0.23	0.62

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Winter injury score based on a scale of 1 to 9 with 9 being the greatest amount of injury.

³ Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

In tables 13, 14, and 15, an open block indicates that the variety was not in that particular test (labeled at the top of the column); an “x” in the block means that the variety was in the test but yielded significantly less than the top-yielding variety. A single asterisk (*) means that the variety was not significantly different from the top variety, based on the 0.05 LSD. It is best to choose a variety that has performed well over several years and locations. Remember to consider the relative spring maturity and the distribu-

tion of yield across the growing season when evaluating productivity of ryegrass varieties (tables 3 through 12).

Tables 16, 17, and 18 are summaries of yield data from 1999 to 2016 of commercial varieties that have been entered in the Kentucky trials. The data are listed as a percentage of the mean of the commercial varieties entered in each specific trial. In other words, the mean for each trial is 100 percent—varieties with percentages over 100 yielded better than average, and varieties with percentages less than 100 yielded lower than average. Direct,

statistical comparisons of varieties cannot be made using the summary tables 16, 17, and 18, but these comparisons do help to identify varieties for further consideration. Varieties that have performed better than average over many years and at several locations have stable performance; others may have performed well in wet years or on particular soil types. These details may influence variety choice and the information can be found in the yearly reports. See the footnotes in tables 16, 17, and 18 to determine to which yearly report to refer.

Table 6. Dry matter yields, seedling vigor, winter injury, plant height, maturity and stand persistence of annual ryegrass varieties sown September 3, 2015 at Lexington, Kentucky (See Table 16 for designation of Italian or Westerwolds type commercial varieties).

Variety	Seedling Vigor ¹ Oct 15, 2015	Percent Stand				Winter Injury ² Jan 29	Plant Height (in) Apr 18	Maturity ³			Yield (tons/acre)						
		2015		2016				2016			2015		2016				
		Oct 15	Mar 18	Jul 5	Aug 5			Apr 18	May 13	Jun 9	Nov 23	Dec 17	Apr 18	May 13	Jun 9	Jul 5	Total
Commercial Varieties-Available for Farm Use																	
Melquatro	4.4	100	100	100	100	2.3	14	31.8	52.0	56.0	0.75	0.45	1.93	2.07	1.32	0.33	6.85*
Barmultra II	3.5	100	100	100	100	2.3	15	32.5	51.0	55.0	0.95	0.57	2.10	1.45	0.93	0.37	6.36*
Nelson	4.8	100	56	45	1	5.5	13	31.8	56.0	56.0	1.10	0.91	0.78	1.42	0.99	0.16	5.35
TetraPrime	1.9	99	100	100	100	0.9	14	32.0	50.0	51.0	0.26	0.42	1.64	1.90	0.59	0.46	5.27
Meroa	4.1	100	99	94	94	3.5	14	32.0	53.5	59.0	0.74	0.57	1.41	1.46	0.81	0.20	5.20
Oryx	4.1	100	100	99	96	4.3	13	31.3	51.0	56.5	0.79	0.52	1.40	1.35	0.74	0.28	5.08
Marshall	3.8	99	99	44	0	1.1	18	32.0	52.5	55.0	0.95	0.54	1.57	1.36	0.59	0.08	5.08
Jackson	3.6	100	97	14	0	4.3	15	32.0	53.5	55.5	0.91	0.68	1.40	1.38	0.61	0.04	5.02
Kowinearly	2.0	100	99	43	0	2.8	16	32.0	56.0	57.0	0.48	0.73	1.59	1.26	0.73	0.09	4.88
Bruiser	4.1	100	97	78	4	4.0	16	32.0	55.5	57.0	0.69	0.71	1.26	1.41	0.68	0.07	4.81
Kospeed	3.3	100	87	15	1	4.5	15	32.5	54.5	57.5	0.92	0.61	1.36	1.24	0.49	0.07	4.68
Feast II	4.1	100	13	48	35	9.0	6	29.0	54.0	58.0	0.55	0.80	0.33	1.36	1.11	0.29	4.45
Fria	3.3	100	90	19	0	5.3	15	31.8	55.5	55.5	0.65	0.72	0.92	1.30	0.70	0.04	4.33
Gulf	4.6	100	6	1	1	9.0	5	29.5	56.5	60.0	0.62	0.77	0.23	0.86	0.54	0.01	3.03
Experimental Varieties																	
BAR LM 15425	1.8	99	100	98	94	1.8	17	32.3	52.0	57.5	0.38	0.61	2.08	1.66	1.06	0.35	6.14*
BAR LM 15426	1.5	95	95	96	68	1.5	17	32.5	52.0	55.5	0.45	0.61	1.73	1.79	0.87	0.30	5.75
BAR LM 15427	1.6	100	100	100	94	0.9	17	32.3	52.5	57.0	0.43	0.62	1.82	1.66	0.83	0.38	5.75
BAR LM 15371	2.1	99	100	100	99	2.5	15	32.0	53.0	55.0	0.47	0.43	1.78	1.50	0.91	0.34	5.44
ME4	3.6	100	100	60	0	2.5	19	32.5	53.0	56.0	0.90	0.52	1.72	1.39	0.53	0.13	5.19
M2CVS	3.5	99	100	61	1	0.8	19	32.5	51.0	56.5	0.77	0.57	1.56	1.30	0.85	0.13	5.18
ME94	5.0	100	100	39	1	3.8	16	32.3	54.5	55.5	0.83	0.69	1.52	1.16	0.73	0.06	4.99
PPG-TAR113	1.3	100	100	90	38	1.8	11	31.3	54.0	57.0	0.07	0.09	0.88	1.42	0.70	0.31	3.48
Mean	3.3	99	88	66	42	3.4	14	32.0	53.0	56.0	0.67	0.60	1.41	1.44	0.79	0.20	5.11
CV,%	15.9	2	9	24	21	37.2	12	2.0	3.0	2.0	36.33	27.67	25.23	18.22	35.21	43.96	14.14
LSD,0.05	0.7	3	11	22	12	1.8	2	1.0	2.0	2.0	0.34	0.23	0.50	0.37	0.39	0.13	1.02

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Winter injury score based on a scale of 1 to 9 with 9 being the greatest amount of injury.

³ Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 7. Dry matter yields, seedling vigor, maturity and stand persistence of perennial ryegrass varieties sown September 5, 2013 at Lexington, Kentucky (See Table 17 for designation of diploid or tetraploid type commercial varieties).

Variety	Seedling Vigor ¹ Oct 14, 2013	Maturity ²										Percent Stand				Yield (tons/acre)						
		2014		2015		2016		2013		2014		2015		2016		2014		2015		2016		3-year Total
		May 7	Jun 12	May 13	May 13	Jun 24	Oct 14	Apr 2	Oct 27	Apr 3	Oct 29	Mar 23	Oct 17	Total	Total	May 14	Jun 24	Total				
Commercial Varieties-Available for Farm Use																						
Kentaur	3.8	32.8	62.0	40.5	45.0	66.0	97	99	99	97	98	96	90	3.80	1.11	1.24	1.21	2.46	7.37*			
Everlast	4.8	56.0	52.8	58.0	58.0	29.0	100	93	98	91	97	96	82	2.87	1.57	1.89	0.25	2.14	6.58*			
Victorian	4.4	56.0	29.0	57.5	58.5	29.0	99	100	100	98	98	98	92	3.02	1.43	1.84	0.25	2.08	6.53*			
PayDay	3.0	34.5	62.0	43.5	47.5	66.0	98	98	100	99	99	98	89	3.41	1.31	1.32	0.44	1.77	6.49*			
Boost	2.3	46.3	61.5	43.5	50.0	64.0	88	88	93	94	85	85	81	3.49	1.35	1.08	0.53	1.61	6.44*			
Calibra	3.5	38.0	62.0	40.5	48.8	65.0	98	98	98	99	98	98	94	3.05	1.19	1.19	0.63	1.81	6.05			
Linn (certified)	3.9	52.0	29.0	55.5	56.0	29.0	99	99	98	97	96	96	66	2.53	1.39	1.71	0.36	2.07	5.98			
Power	3.8	35.8	61.5	40.5	49.3	66.0	98	99	100	97	98	98	92	3.36	1.20	0.89	0.52	1.40	5.97			
BG34	3.8	32.3	53.8	39.0	45.0	64.0	99	100	100	88	91	91	79	3.11	0.91	0.78	0.49	1.27	5.29			
Experimental Varieties																						
RAD-MFP145	2.9	33.8	53.8	43.5	50.0	66.0	97	97	98	99	97	97	90	3.43	1.56	1.20	0.91	2.11	7.10*			
RAD-MFP141	2.9	32.8	61.5	39.0	47.5	66.0	97	97	99	97	98	97	91	3.63	1.41	1.06	0.72	1.78	6.83*			
Mean	3.5	40.9	53.5	45.5	50.5	55.5	97	97	98	96	98	95	86	3.25	1.31	1.29	0.57	1.86	6.42			
CV,%	13.0	9.4	16.5	4.7	3.7	2.1	3	3	2	3	6	6	12	10.27	19.88	34.22	38.09	25.78	11.55			
LSD,0.05	0.7	5.6	12.7	3.1	2.7	1.7	4	4	3	5	8	8	15	0.48	0.38	0.64	0.32	0.69	1.07			

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 8. Dry matter yields, seedling vigor, winter injury, maturity, and stand persistence of perennial ryegrass varieties sown September 5, 2014, at Lexington, Kentucky (See Table 17 for designation of diploid or tetraploid type commercial varieties).

Variety	Seedling Vigor ¹ Oct 9, 2014	Winter Injury ² Jan 22, 2015	Maturity ³			Percent Stand					Yield (tons/acre)			2-year Total	
			2015		2016	2014	2015		2016		2015	2016			
			May 11	May 13	Jun 24	Oct 9	Apr 3	Oct 29	Mar 21	Oct 17	Total	May 13	Jun 24		Total
Commercial Varieties-Available for Farm Use															
TetraMag	4.6	6.8	47.5	47.5	62.0	96	94	95	96	66	5.45	2.34	1.42	3.76	9.21*
Remington PLUS NEA2	3.8	3.0	45.0	45.0	62.0	98	99	99	99	95	4.60	1.94	1.52	3.46	8.05*
Grand Daddy	2.6	2.0	56.0	57.5	53.8	95	96	96	96	75	4.52	2.16	0.59	2.75	7.27
Albion	2.3	3.0	46.3	45.0	53.8	94	98	97	95	84	4.40	1.67	1.06	2.72	7.13
Power	4.0	5.5	49.3	52.5	37.3	95	96	97	97	76	4.09	1.91	0.82	2.72	6.81
PayDay	3.3	3.5	46.3	51.5	45.5	96	97	98	98	83	4.08	1.78	0.84	2.62	6.70
Remington	2.6	1.3	46.3	46.3	62.0	94	97	97	98	86	3.97	1.26	1.22	2.48	6.44
Calibra	4.3	3.3	46.3	50.0	53.8	94	95	97	97	84	3.59	1.70	0.60	2.30	5.89
BG34	3.8	2.3	47.5	46.3	45.5	99	99	99	96	70	3.44	1.45	0.83	2.28	5.73
Victorian	3.8	8.5	56.0	56.0	37.3	100	97	97	96	61	3.52	1.71	0.42	2.13	5.65
Linn (certified)	3.5	5.5	56.0	56.0	29.0	98	98	98	98	70	3.49	1.77	0.38	2.14	5.63
Experimental Varieties															
GO-AX11	3.8	6.0	47.5	39.3	62.0	95	91	93	93	33	4.55	2.43	1.18	3.61	8.15*
13BSTRD	3.3	2.5	49.8	52.0	53.8	95	96	96	96	66	4.16	2.36	0.67	3.03	7.19
13BSTYW	3.6	2.8	47.5	51.5	62.0	95	97	90	80	54	4.19	1.92	0.92	2.83	7.03
AGR1P-156 AR1	3.9	3.8	53.5	53.5	29.0	98	99	99	99	58	3.95	2.03	0.83	2.85	6.80
GO-13ALF	2.6	3.0	46.3	45.0	45.5	96	99	99	98	86	4.28	1.29	1.22	2.50	6.79
13PI2B1	3.6	4.3	50.8	49.3	37.3	100	100	98	98	66	4.43	1.53	0.58	2.11	6.54
13PI2B2	3.8	2.3	49.0	50.3	37.3	100	100	99	98	61	3.77	1.90	0.70	2.60	6.37
13PI3B	3.9	2.3	51.5	48.0	53.8	100	100	100	99	64	3.89	1.55	0.68	2.23	6.12
GO-13ABFR	2.9	2.8	45.0	45.0	62.0	96	97	97	95	86	3.70	1.27	1.09	2.37	6.07
GO-13AXT	3.1	2.3	49.3	48.5	53.8	97	99	97	86	60	3.70	1.49	0.87	2.36	6.06
AGR1P-157 AR1	4.0	2.3	45.0	47.5	45.5	100	100	100	99	89	3.50	1.51	0.87	2.38	5.88
Mean	3.5	3.6	49.0	49.2	49.3	97	97	97	96	71	4.06	1.77	0.88	2.65	6.70
CV,%	18.3	36.6	4.8	10.9	28.5	3	2	3	9	29	11.51	24.30	36.64	21.55	13.15
LSD,0.05	0.9	1.8	3.3	7.6	19.8	4	3	5	12	29	0.66	0.61	0.45	0.81	1.25

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Winter injury score based on a scale of 1 to 9 with 9 being the greatest amount of injury.

³ Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Summary

Selecting a good variety of annual or perennial ryegrass or festulolium is an important first step in establishing a productive stand of grass. Proper management, beginning with seedbed preparation and continuing throughout the life of the stand, is necessary for even the highest-yielding variety to produce to its genetic potential.

The following is a list of University of Kentucky Cooperative Extension publications related to ryegrass management.

They are available from your county Extension office and are listed in the “Publications” section of the UK Forage website, www.uky.edu/Ag/Forage.

- Lime and Fertilizer Recommendations (AGR-1)
- Grain and Forage Crop Guide for Kentucky (AGR-18)
- Establishing Forage Crops (AGR-64)
- Forage Identification and Use Guide (AGR-175)
- Annual Ryegrass (AGR-179)
- New Recommendations for Perennial Ryegrass Seedings for Kentucky Horse Farms (ID-142)

- Rotational Grazing (ID-143)
- Establishing and Managing Horse Pastures (ID-147)
- Festulolium Hybrid Grass (see the UK Forage website under publications and grasses)

Authors

G.L. Olson is a research specialist and S.R. Smith is an Extension professor of Forages. T.D. Phillips is an associate professor of Tall Fescue Breeding.

Table 9. Dry matter yields, seedling vigor, winter injury, maturity, and stand persistence of perennial ryegrass varieties sown September 3, 2015, at Lexington, Kentucky (see Table 17 for designation of diploid or tetraploid type commercial varieties).

Variety	Seedling Vigor ¹ Oct 15, 2015	Winter Injury ² Jan 29, 2016	Maturity ³		Percent Stand			Yield (tons/acre)		
			2016		2015	2016		2016		
			May 13	Jun 24	Oct 15	Mar 18	Oct 17	May 13	Jun 24	Total
Commercial Varieties-Available for Farm Use										
Barvitra	5.0	2.0	55.5	62.0	100	100	35	4.12	1.48	5.60*
Grand Daddy	2.9	1.1	55.0	62.0	99	100	76	4.05	1.41	5.47*
Remington	3.3	0.6	45.0	29.0	100	100	100	3.76	0.91	4.67
Power	3.6	1.3	53.5	29.0	100	100	96	3.89	0.74	4.62
Albion	2.9	0.9	46.3	29.0	86	99	95	3.79	0.81	4.60
Calibra	4.1	0.6	51.5	29.0	100	100	94	3.82	0.77	4.59
Remington PLUS NEA2	2.9	0.6	46.8	29.0	100	100	100	3.45	0.98	4.43
Linn (certified)	3.4	3.0	56.0	29.0	100	100	73	3.27	0.52	3.79
BG-34	3.8	1.0	49.8	29.0	100	100	90	2.85	0.70	3.54
Experimental Varieties										
BAR LP 15261	2.5	0.8	47.5	37.3	100	100	100	3.49	0.86	4.35
GPT-14021	2.5	0.8	48.0	37.3	99	100	100	3.32	0.85	4.18
GPD-14017	3.1	2.5	55.0	37.3	100	100	79	3.36	0.61	3.97
GPD-14018	3.3	1.8	48.0	29.0	100	100	97	2.88	0.68	3.56
GPT-14023	3.0	6.0	55.5	62.0	100	55	70	2.52	0.95	3.47
TAL-PR-04	3.9	4.5	56.0	62.0	100	85	71	2.83	0.42	3.25
TAL-PR-02	3.8	6.0	56.0	62.0	100	74	69	2.77	0.30	3.07
GPD-14019	2.8	6.8	50.8	62.0	99	40	68	1.92	0.83	2.75
TAL-PR-03	1.8	4.3	56.0	37.3	100	97	30	2.15	0.31	2.46
Mean	3.2	2.5	51.8	41.9	99	92	80	3.23	0.79	4.02
CV,%	17.3	26.3	4.1	18.8	1	10	22	11.97	45.51	13.59
LSD,0.05	0.8	0.9	3.0	11.1	2	13	25	0.55	0.51	0.78

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Winter injury score based on a scale of 1 to 9 with 9 being the greatest amount of injury.

³ Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 10. Dry matter yields, seedling vigor, maturity, and stand persistence of festulolium varieties sown September 5, 2013, at Lexington, Kentucky.

Variety	Seedling Vigor ¹ Oct 14, 2013	Maturity ²					Percent Stand							Yield (tons/acre)					3-year Total
		2014		2015		2016	2013	2014		2015		2016		2014	2015	2016			
		May 8	Jun 12	May 11	May 13	Jun 21	Oct 14	Apr 2	Oct 28	Apr 3	Oct 29	Mar 23	Oct 17	Total	Total	May 13	Jun 23	Total	
Commercial Varieties-Available for Farm Use																			
Hykor	2.9	54.5	29.0	57.0	57.5	29.0	100	93	99	99	98	97	97	4.00	2.67	1.65	0.78	2.43	9.10*
Perun	4.0	41.0	62.0	45.0	52.0	62.0	99	99	98	85	84	84	48	5.36	1.24	1.75	0.75	2.50	9.10*
Lofa	4.1	41.0	62.0	45.0	50.5	62.0	99	99	99	91	89	88	43	4.99	1.66	1.41	0.89	2.30	8.95*
Perseus	3.9	42.3	62.0	45.0	50.5	62.0	100	99	99	86	88	88	56	5.02	1.65	1.36	0.56	1.92	8.60*
Barfest	3.3	43.0	62.0	45.0	50.0	62.0	99	98	99	98	95	94	60	4.52	1.56	1.55	0.68	2.23	8.31*
Felina	2.4	54.0	29.0	56.5	57.0	29.0	97	89	98	97	96	94	93	3.58	2.13	1.53	0.68	2.21	7.93*
Hostyn	3.0	47.3	62.0	45.0	52.5	61.5	96	94	95	84	69	69	35	4.73	1.24	1.27	0.43	1.70	7.67
Spring Green	3.5	42.3	62.0	45.0	50.0	62.0	100	100	100	99	97	97	74	4.34	1.43	1.44	0.45	1.88	7.65
Mahulena	1.8	56.0	29.0	58.0	59.5	29.0	98	84	95	97	95	95	94	3.21	2.07	1.52	0.80	2.32	7.60
Duo	3.5	46.8	62.0	45.0	51.0	62.0	100	100	100	99	96	94	63	3.88	1.23	1.22	0.36	1.58	6.70
Rebab	2.3	50.3	43.5	54.5	55.5	29.0	97	92	97	97	97	94	94	3.01	2.07	1.06	0.43	1.49	6.56
Fojtan	2.0	48.5	50.8	54.5	56.5	29.0	98	85	96	97	95	96	95	3.00	1.88	0.84	0.67	1.51	6.40
Gain	4.8	53.5	61.0	45.0	50.0	62.0	100	76	80	64	53	43	15	3.73	1.10	0.21	0.21	0.42	5.26
Bonus	4.8	51.5	62.0	48.7	—	—	100	6	4	4	4	6	0	1.94	0.45	0.00	0.00	0.00	2.39
Meadow Green	5.0	47.5	61.5	52.0	—	—	100	18	4	1	1	1	0	2.18	0.18	0.00	0.00	0.00	2.35
Experimental Varieties																			
KYFA9819	2.6	40.3	62.0	47.8	53.0	62.0	100	97	97	92	90	91	48	4.06	1.46	1.83	0.54	2.36	7.88*
Amp1427	2.3	34.5	62.0	45.0	50.5	61.5	99	95	97	97	95	96	62	3.92	1.37	1.18	0.67	1.85	7.13
XLFFL	4.3	52.5	61.5	54.0	—	—	100	71	13	4	3	3	0	3.22	0.21	0.00	0.00	0.00	3.43
Mean	3.3	47.0	54.7	49.1	53.2	50.4	99	83	82	77	75	74	54	3.82	1.42	1.10	0.49	1.60	6.83
CV,%	17.4	8.9	9.5	3.7	2.8	0.8	2	9	2	5	8	10	30	11.32	18.40	40.25	52.03	32.86	14.03
LSD,0.05	0.8	6.0	7.3	2.9	2.4	0.6	2	11	3	6	9	10	23	0.61	0.37	0.63	0.37	0.74	1.36

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 11. Dry matter yields, seedling vigor, maturity, and stand persistence of festulolium varieties sown September 5, 2014, at Lexington, Kentucky.

Variety	Seedling Vigor ¹ Oct 9, 2014	Maturity ²			Percent Stand					Yield (tons/acre)				2-year Total
		2015		2016	2014	2015		2016		2015	2016			
		May 6	May 13	Jun 21	Oct 9	Apr 2	Oct 29	Mar 21	Oct 17	Total	May 13	Jun 21	Total	
Commercial Varieties-Available for Farm Use														
Hykor	3.5	52.0	59.5	29.0	98	97	98	98	97	3.65	1.58	0.77	2.35	6.00*
Lofa	5.0	33.0	52.0	62.0	100	100	98	98	45	3.61	1.43	0.62	2.06	5.67*
Perseus	4.8	32.8	49.3	62.0	100	100	100	98	35	3.57	1.03	0.97	2.00	5.57*
Perun	3.9	32.8	53.5	62.0	99	100	99	98	30	3.63	1.18	0.76	1.94	5.57*
Mahulena	2.0	56.0	60.0	29.0	70	87	90	93	93	3.16	1.45	0.80	2.24	5.40*
Hostyn	4.0	32.8	54.0	62.0	96	97	97	94	30	3.40	1.18	0.79	1.97	5.37*
Spring Green	4.3	40.5	52.0	62.0	98	99	99	98	84	3.27	1.26	0.67	1.94	5.20*
Felina	2.3	47.5	60.0	29.0	89	94	96	96	96	2.71	1.17	0.95	2.12	4.83*
Duo	3.8	49.3	51.0	62.0	96	98	98	98	75	3.22	1.00	0.62	1.61	4.83*
Barfest	4.5	33.8	47.5	62.0	98	99	99	98	76	3.09	0.96	0.56	1.52	4.60
Rebab	2.3	40.5	56.5	29.0	89	92	93	95	95	2.33	0.92	0.64	1.56	3.89
Fojtan	2.0	39.0	56.5	29.0	85	94	95	95	94	1.90	1.14	0.60	1.74	3.64
Experimental Varieties														
KYFA9819	4.3	32.5	49.8	62.0	98	97	98	97	50	3.01	1.18	0.66	1.83	4.84*
PPG-FEST 102	3.4	39.0	49.8	62.0	93	96	96	95	55	2.89	0.73	0.78	1.50	4.40
Mean	3.6	40.1	53.7	50.2	93	96	97	96	68	3.10	1.16	0.73	1.88	1.99
CV,%	15.1	6.5	3.4	0.0	6	2	2	2	15	17.54	33.21	31.67	25.13	18.40
LSD,0.05	0.8	3.7	2.6	0.0	8	3	2	3	15	0.78	0.55	0.33	0.68	1.31

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 12. Dry matter yields, seedling vigor, winter injury, maturity, and stand persistence of festulolium varieties sown September 3, 2015, at Lexington, Kentucky.

Variety	Seedling Vigor ¹ Oct 15, 2015	Winter Injury ² Jan 29, 2016	Maturity ³		Percent Stand			Yield (tons/acre)				
			2016		2015	2016		2015		2016		
			May 11	Jun 20	Oct 15	Mar 18	Oct 17	Nov 23	Dec 18	May 12	Jun 20	Total
Commercial Varieties-Available for Farm Use												
Perseus	3.4	1.3	50.0	56.5	100	100	50	0.51	0.51	5.32	1.58	7.91*
Perun	4.1	1.6	51.0	56.5	100	100	38	0.42	0.55	5.45	1.25	7.68*
Spring Green	3.0	1.0	51.5	57.0	100	100	95	0.60	0.58	4.87	1.04	7.09*
Lofa	4.3	1.3	50.5	56.0	100	100	74	0.30	0.30	4.45	1.22	6.25*
Barfest	2.9	0.9	50.5	40.5	100	100	95	0.21	0.49	4.25	0.90	5.86
Duo	4.3	4.3	56.0	60.0	100	96	96	0.46	0.79	3.62	0.94	5.81
Hykor	1.5	0.5	57.0	29.0	100	100	100	0.12	0.35	3.52	1.44	5.44
Fojtan	1.3	0.5	56.0	29.0	100	100	100	0.02	0.10	3.22	1.12	4.46
Experimental Varieties												
KYFL1013	4.1	1.3	50.0	53.5	100	100	98	0.35	0.53	5.33	1.04	7.24*
PPG-FEST-102	2.4	1.0	53.5	57.0	100	100	97	0.35	0.46	4.18	0.86	5.85
Mean	3.1	1.4	52.6	49.5	100	100	84	0.33	0.47	4.42	1.14	6.36
CV,%	15.9	31.5	1.8	9.0	0	0	23	67.19	66.06	19.03	23.76	21.06
LSD,0.05	0.7	0.6	1.4	6.5	1	1	28	0.33	0.45	1.22	0.39	1.94

¹ Vigor score based on a scale of 1 to 5 with 5 being the most vigorous seedling growth.

² Winter injury score based on a scale of 1 to 9 with 9 being the greatest amount of injury.

³ Maturity rating scale: 37 = flag leaf emergence, 45 = boot swollen, 50 = beginning of inflorescence emergence, 58 = complete emergence of inflorescence, 62 = beginning of pollen shed. See Table 2 for complete scale.

*Not significantly different from the highest numerical value in the column, based on the 0.05 LSD.

Table 13. Performance of annual ryegrass varieties sown in 2015 at Lexington.¹

Variety	Type	Proprietor/KY Distributor	2015 ²
			2016 ³
Commercial Varieties-Available for Farm Use			
Barmultra II	Italian tetraploid	Barenbrug USA	*
Bruiser	Westerwold diploid	Ampac Seed	x ⁴
Feast II	Italian tetraploid		x
Fria	Westerwold diploid	Allied Seed	x
Gulf		Public	x
Jackson		The Wax Company	x
Kospeed		Smith Seed	x
Kowinearly		Services	x
Marshall		The Wax Company	x
Melquatro		Italian tetraploid	Hood River Seed
Meroa	Westerwold diploid	Smith Seed Services	x
Nelson	Westerwold tetraploid	The Wax Company	x
Oryx	Italian diploid	Hood River Seed	x
TetraPrime	Italian tetraploid	Mountain View Seeds	x
Experimental Varieties			
BAR LM 15425	Westerwold tetraploid	Barenbrug USA	*
BAR LM 15426			x
BAR LM 15427			x
BAR LM 15371			x
ME4	Westerwold diploid	The Wax Company	x
ME-94			x
M2CVS			x
PPG-TAR113			Smith Seed Services

¹ See Table 16 for summary of yield data on named varieties from 2000-2016.

² Establishment year.

³ Harvest year.

⁴ "x" in the box indicates the variety was in the test but yielded significantly less than the top yielding variety.

*Not significantly different from the highest yielding variety in the test.

Table 14. Performance of perennial ryegrass across years at Lexington.

Variety	Type	Proprietor/KY Distributor	2013 ¹			2014		2015
			2014 ²	2015	2016	2015	2016	2016
Commercial Varieties-Available for Farm Use								
Albion	tetraploid	Grassland Oregon				x ³	x	x
Barvitra	hybrid	Barenbrug USA						*
BG34	diploid	Barenbrug USA	x	x	x	x	x	x
Boost	tetraploid	Allied Seed	*	*	x			
Calibra	tetraploid	DLF International	x	*	*	x	x	x
Everlast	diploid	Caudill Seed	x	*	*			
Grand Daddy	tetraploid	Smith Seed Services				x	x	*
Kentaur	tetraploid	DLF International	*	x	*			
Linn (certified)	diploid	Public	x	*	*	x	x	x
PayDay	tetraploid	Mountain View Seeds	*	*	*	x	x	
Power	tetraploid	Ampac Seed Company	*	*	x	x	x	x
Remington	tetraploid	Barenbrug USA				x	x	x
Remington PLUS NEA2	tetraploid	Barenbrug USA				x	*	x
TetraMag	tetraploid	Mountain View Seeds				*	*	
Victorian	diploid	Caudill Seed	x	*	*	x	x	
Experimental Varieties								
AGRLP-156AR1	—	Ag. Research				x	x	
AGRLP-157AR1	—	Ag. Research				x	x	
BAR LP 15261	tetraploid	Barenbrug USA						x
GO-AX11	intermediate	Grassland Oregon				x	*	
GO-13ABFR	tetraploid	Grassland Oregon				x	x	
GO-13ALF	tetraploid	Grassland Oregon				x	x	
GO-13AXT	tetraploid	Grassland Oregon				x	x	
GPD-14017AR1	diploid	Ag. Research						x
GPD-14018AR1	diploid	Ag. Research						x
GPD-14019	diploid	Ag. Research						x
GPT-14021	tetraploid	Ag. Research						x
GPT-14023AR5	tetraploid	Ag. Research						x
RAD-MFP141	tetraploid	Radix Research	*	*	*			
RAD-MFP145	tetraploid	Radix Research	*	*	*			
TAL-PR-02	diploid	Ag. Research						x
TAL-PR-03	diploid	Ag. Research						x
TAL-PR-04	diploid	Ag. Research						x
13BSTRD	—	Oregro Seeds				x	*	
13BSTYW	—	Oregro Seeds				x	x	
13P12B1	—	Oregro Seeds				x	x	
13P12B2	—	Oregro Seeds				x	x	
13P13B	—	Oregro Seeds				x	x	

¹ Establishment year.

² Harvest year.

³ "x" in the box indicates the variety was in the test but yielded significantly less than the top yielding variety. Open boxes indicate the variety was not in the test.

*Not significantly different from the highest yielding variety in the test.

Table 15. Performance of festulolium varieties across years at Lexington.

Variety	Type ²	Proprietor/KY Distributor	2013 ¹			2014		2015
			2014 ³	2015	2016	2015	2016	2016
Commercial Varieties-Available for Farm Use								
Barfest	MF x PR	Barenbrug USA	x ⁴	x	*	*	x	x
Duo	MF x PR	Ampac Seed	x	x	x	*	x	x
Felina	(TF x IR) x TF	DLF International	x	x	*	x	*	
Gain	MF x IR	Allied Seed	x	x	x			
Fojtan	(TF x IR) x TF	DLF International	x	x	x	x	*	x
Hostyn	MF x IR	DLF International	x	x	*	*	*	
Hykor	(TF x IR) x TF	DLF International	x	*	*	*	*	x
Lofa	(TF x Int) x Int	DLF International	*	x	*	*	*	*
Mahulena	(TF x IR) x TF	DLF International	x	x	*	*	*	
Meadow Green	—	Pure Seed	x	x	x			
Perseus	MF x IR	DLF International	*	x	*	*	*	*
Perun	MF x IR	DLF International	*	x	*	*	*	*
Rebab	(TF x IR) x TF	DLF International	x	x	x	x	x	
Spring Green	MF x PR	Turf Seed	x	x	*	*	*	*
Bonus	MF x IR	Allied Seed	x	x	x			
Experimental Varieties								
Amp1427	—	Ampac Seed	x	x	*			
KYFL1013	MF x IR	KY Agric. Exp. Station						*
KYFA9819	MF x IR	KY Agric. Exp. Station	x	x	*	*	*	
PPG-FEST-102	PR x MF	Mountain View Seeds				*	x	x
XLF FL	—	ProSeeds Marketing	x	x	x			

¹ Establishment year.

² MF = meadow fescue, TF = tall fescue, IR = Italian ryegrass, PR = perennial ryegrass, Int = intermediate ryegrass

³ Harvest year.

⁴ "x" in the box indicates the variety was in the test but yielded significantly less than the top yielding variety. Open boxes indicate the variety was not in the test.

*Not significantly different from the highest yielding variety in the test.

Table 18. Summary of Kentucky festulolium yield trials 2001-2016 (yield shown as a percentage of the mean of the commercial varieties in the trial).¹

Variety	Type ²	Proprietor	Lexington									Mean ⁵ (#trials)
			2001 ^{3,4}	2005	2008	2009	2010	2011	2012	2013	2014	
			2yr ⁶	3yr	3yr	3yr	3yr	3yr	2yr	3yr	2yr	
Agula	MF x IR	Allied Seed					94			–		
Barfest	MF x PR	Barenbrug USA					105	101	107	119	91	105(5)
Bonus	MF x IR	Allied Seed					93	46	32	34		51(4)
Duo	MF x PR	Ampac Seed		89	98	99	95	106	103	96	96	98(8)
Felina	(TF x IR) x TF	DLF International	104				132	118	134	114	96	116(6)
Fojtan	(TF x IR) x TF	DLF International					112	101	124	92	72	100(5)
Gain	MF x IR	Allied Seed					103	77	52	75		77(4)
Hostyn	MF x IR	DLF International							107	110	106	108(3)
Hykor	(TF x IR) x TF	DLF International					133	141	153	131	119	135(5)
Lofa	(TF x Int) x Int	DLF International					105	107	110	128	112	112(5)
Mahulena	(TF x IR) x TF	DLF International							131	109	107	116(3)
Meadow Green	–	Pure Seed							37	34		36(2)
Perseus	MF x IR	DLF International					132	114	126	123	110	121(5)
Perun	MF x IR	DLF International					127	114	107	131	110	118(5)
Rebab	(TFxIR) xTF	DLF International								94	77	86(2)
Spring Green	MF x PR	Turf-Seed	96	111	114	101	113	112	114	110	103	108(9)
Sweet Tart	MF x IR	ProSeeds Marketing			88		82	63	62			74(4)

¹ The festuloliums were in fescue trials from 2001-2005 and in perennial ryegrass trials from 2008-2009.

² MF = meadow fescue, TF = tall fescue, IR = Italian ryegrass, PR = perennial ryegrass, Int = intermediate ryegrass.

³ Year trial was established.

⁴ Use this summary table as a guide in making variety decisions, but refer to specific yearly reports to determine statistical differences in forage yield between varieties.

To find actual yields, look in the yearly report for the final year of each specific trial. For example, the Lexington trial planted in 2012 was harvested three years, so the final report would be “2015 Annual and Perennial Ryegrass and Festulolium Report” archived in the KY Forage website at <www.uky.edu/Ag/Forage>.

⁵ Mean only presented when respective variety was included in two or more trials.

⁶ Number of years of data.